

Sedación consciente.

Elección conveniente.

Velocidad de infusión en ml/hr	DOSIS (µg/kg)					
	0.2	0.3	0.4	0.5	0.6	0.7
40	2.0	3.0	4.0	5.0	6.0	7.0
45	2.3	3.4	4.5	5.6	6.8	7.9
50	2.5	3.8	5.0	6.3	7.5	8.8
55	2.8	4.1	5.5	6.9	8.3	9.6
60	3.0	4.5	6.0	7.5	9.0	10.5
65	3.3	4.9	6.5	8.1	9.8	11.4
70	3.5	5.3	7.0	8.8	10.5	12.3
75	3.8	5.6	7.5	9.4	11.3	13.1
80	4.0	6.0	8.0	10.0	12.0	14.0
85	4.3	6.4	8.5	10.6	12.8	14.9
90	4.5	6.8	9.0	11.3	13.5	15.8
95	4.8	7.1	9.5	11.9	14.3	16.6
100	5.0	7.5	10.0	12.5	15.0	17.5
105	5.3	7.9	10.5	13.1	15.8	18.4
110	5.5	8.3	11.0	13.8	16.5	19.3
115	5.8	8.6	11.5	14.4	17.3	20.1
120	6.0	9.0	12.0	15.0	18.0	21.0
125	6.3	9.4	12.5	15.6	18.8	21.9
130	6.5	9.8	13.0	16.3	19.5	22.8
135	6.8	10.1	13.5	16.9	20.3	23.6
140	7.0	10.5	14.0	17.5	21.0	24.5
145	7.3	10.9	14.5	18.1	21.8	25.4
150	7.5	11.3	15.0	18.8	22.5	26.3

Peso  
(Kg)

Uso de la Tabla de Dosificación

- 1\_ Peso del paciente (Filas)
- 2\_ Dosis que desea administrar (Columnas)
- 3\_ Velocidad de infusión (Intersección).

<http://www.learnicu.org/SiteCollectionDocuments/Pain,%20Agitation,%20Delirium.pdf>  
[http://scielo.isciii.es/ing/revistas/neuro/v16ns/2\\_t2.jpg](http://scielo.isciii.es/ing/revistas/neuro/v16ns/2_t2.jpg)